


- (6) 1. Given the graph of f below, determine each of the following. Use ∞ , $-\infty$ or “does not exist” (DNE) where appropriate.



(a) $\lim_{x \rightarrow -2} f(x) =$

(b) $\lim_{x \rightarrow -3^+} f(x) =$

(c) $f(-2) =$

(d) $f'(0) =$

(e) $\lim_{x \rightarrow \infty} f(x) =$

(f) $\lim_{x \rightarrow 2^-} f(x) =$

(g) $\lim_{x \rightarrow -\infty} f(x) =$

(h) $\lim_{x \rightarrow 2} f(x) =$

(i) $f'(1) =$

(j) List all x -values where the function is discontinuous and name the types of discontinuity.

(k) List all x -values where the function is continuous but not differentiable. Justify your answer.

- (21) 2. Evaluate the following limits. Use ∞ , $-\infty$, and dne as appropriate.

(a) $\lim_{x \rightarrow 3} \frac{\frac{3}{7} - \frac{x}{3x-2}}{x-3}$

(b) $\lim_{x \rightarrow 4} \frac{\sqrt{3x} - \sqrt{4x-4}}{2x-8}$

(c) $\lim_{x \rightarrow 10} \frac{x^2 + x - 110}{x + 11}$

(d) $\lim_{x \rightarrow 1} \frac{5x^2 - 5}{15x^2 - 18x + 3}$

(e) $\lim_{x \rightarrow -4} \frac{x^2 - 3x - 28}{(x+4)^3}$

(f) $\lim_{x \rightarrow -\infty} \frac{(x^2 - 3x^3)^2(2x+1)}{(2x^2+1)^2(3x+4)^3}$

(g) $\lim_{x \rightarrow 2^+} \frac{|x-2|}{2x^3 - 4x^2 + 3x - 6}$

- (2) **3.** Answer the following questions with True or False. Justify your answer.
- (a) A function can have AT MOST two horizontal asymptotes.
- (b) A function cannot intersect its vertical asymptote.
- (3) **4.** Use the definition of continuity to determine the points of discontinuity of the following function, $f(x)$. Name the type of discontinuity.

$$f(x) = \begin{cases} \frac{3x + 12}{x^2 + 2x - 8} & : x \leq -2 \\ x^2 & : -2 < x < 2 \\ \frac{4}{3 - x} & : x \geq 2 \end{cases}$$

- (3) **5.** Find the value(s) of k for which the function $g(x)$ is continuous on \mathbb{R} .

$$g(x) = \begin{cases} k^2 + 2x & : x < -1 \\ -kx & : x \geq -1 \end{cases}$$

- (3) **6.**

- (a) State the limit definition of the derivative.
- (b) Use this definition to calculate the derivative of $g(x) = \sqrt{3 - 2x}$

- (21) **7.** Find the derivative of each of the following functions. Do not simplify your answers.

(a) $y = \frac{3}{2x^6} - \sqrt[3]{x^4} + \log_8 x + \pi$

(b) $g(x) = \cot(xe^x)$

(c) $f(x) = \sin^2(x^3 - 7^x)$

(d) $y = \frac{e^{9x} + 1}{\ln(6x^3 - 3x^5) - 3}$

(e) $y = \sqrt[3]{\sec(2x^3 + 4)} + 6$

(f) $(x + 3y)^2 = x^2 + e^{2y}$

(g) $y = (\cos x + 4x)^{\sqrt{x}}$

- (3) **8.** Use logarithmic differentiation to find the derivative of $f(x) = \frac{5(2^x - 6x^3)^5 \ln x}{7(\tan x)^4 \sqrt{(3x^2 + 2x + 1)^5}}$

- (4) **9.** Given $\frac{-4x^2}{y} = x \ln y - y$


- (a) Find y'
(b) Find the equation of the tangent line at the point $(x, y) = (\frac{1}{2}, 1)$
- (3) 10. Find both y' and y'' for $y^2 = xy + 8$
- (4) 11. Find the value(s) where the tangent line to $g(x) = (x^2 + 4)^3(x^2 - 10)^4$ is horizontal.
- (3) 12. Find all critical numbers of the function $h(x) = x\sqrt{1+x}$.
- (3) 13. Find the absolute extrema of the function $g(x) = \frac{x^3 + 4}{x^2}$ on the interval $[1, 4]$

- (10) 14. Consider

$$f(x) = \frac{x}{(2x+1)^2}, f'(x) = \frac{-2x+1}{(2x+1)^3}, \text{ and } f''(x) = \frac{8(x-1)}{(2x+1)^4}$$

Determine the following then neatly sketch a graph of $f(x)$ on the following page.

- (a) all x - and y - intercepts
(b) all vertical and horizontal asymptotes
(c) the intervals on which $f(x)$ is increasing and decreasing
(d) all local (relative) maxima and minima
(e) the intervals on which $f(x)$ is concave up and concave down
(f) any points of inflection
(g) sketch the curve $y = f(x)$ on the following page
- (3) 15. Canadian Apparel prints orange T-shirts featuring the basic differentiation rules. It costs \$2000 to set up the printer and \$12 to produce and print a T-shirt, that is, the cost function of producing x T-shirts is $C(x) = 2000 + 12x$.
- (a) Find average cost function.
(b) What happens to the average cost as the number of T-shirts gets large (that is, as $x \rightarrow \infty$)?
(c) Find the marginal average cost at the production level of 300 T-shirts and interpret the result.
- (4) 16. A group of sadistic calculus teachers decided to lock up some adorable unicorns in an unusual room, because reasons. The room is a rectangle with a isosceles triangle on top, where the base of the rectangle (and triangle) is $8x$ and the sides of the triangle are $5x$.



What is the maximum area for the room, if the total length of the walls of the room add up to 2640m?

- (4) **17.** The demand for a mini samurai robot unicorn is given by $p^2 + 4x = 72081$.
- Find the elasticity of demand function, η , in terms of p .
 - By calculating the value of η when the price is $p = \$9$ (and $x = 18000$), determine what will happen to quantity demanded if the price is increased by 6%.
 - Does the revenue increase or decrease if the price increase in (b) is approved?

Answers

- a) -1 b) 1 c) -3 d) 2 e) -2 f) $-\infty$ g) 2 h) dne i) dne
 j) -3 : jump, -2 : removable, 2 : infinite
 k) -1, 1 : corner, abrupt change in slope
- a) $\frac{2}{49}$ b) $\frac{-1}{4\sqrt{12}} = \frac{-1}{8\sqrt{3}}$ c) 0 d) $\frac{5}{6}$ e) $-\infty$ f) $\frac{1}{6}$ g) $\frac{1}{11}$
- a) True. There can be one on the left as x approaches $-\infty$ and one on the right at x approaches ∞ .
 b) False. It is possible for there to be exactly one point ON the vertical asymptote, but the function cannot intersect it.
- Removable discontinuity at $x = -4$, infinite discontinuity at $x = 3$, and a jump discontinuity at $x = -2$.
- $k = -1, 2$
- (a) $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
 (b) $f'(x) = \frac{-1}{\sqrt{3-2x}}$
- (a) $y' = -9x^{-7} - \frac{4}{3}x^{1/3} + \frac{1}{x \ln 8}$
 (b) $g'(x) = -\csc^2(xe^x)(e^x + xe^x)$
 (c) $f'(x) = 2 \sin(x^3 - 7^x) \cos(x^3 - 7^x)(3x^2 - 7^x \ln 7)$
 (d) $y' = \frac{(9e^{9x})[\ln(6x^3 - 3x^5) - 3] - (e^{9x} + 1) \left(\frac{18x^2 - 15x^4}{6x^3 - 3x^5} \right)}{[\ln(6x^3 - 3x^5) - 3]^2}$
 (e) $y' = \frac{1}{3}[\sec(2x^3 + 4) + 6]^{-2/3} \sec(2x^3 + 4) \tan(2x^3 + 4)(6x^2)$
 (f) $y' = \frac{2x - 2(x + 3y)}{6(x + 3y) - 2e^{2y}} = \frac{-3y}{3x + 9y - e^{2y}}$
 (g) $y' = (\cos x + 4x)^{\sqrt{x}} \left(\frac{1}{2\sqrt{x}} \ln(\cos x + 4x) + \sqrt{x} \left(\frac{-\sin x + 4}{\cos x + 4x} \right) \right)$
- $f'(x) = \left(\frac{5(2^x - 6x^3)^5 \ln x}{7(\tan x)^4 \sqrt{(3x^2 + 2x + 1)^5}} \right) \left(\frac{5(2^x \ln 2 - 18x^2)}{2^x - 6x^3} + \frac{1}{x \ln x} - \frac{\sec^2 x}{\tan x} - \frac{5(6x + 2)}{4(3x^2 + 2x + 1)} \right)$
- a) $y' = \frac{y^2 \ln y + 8xy}{4x^2 - xy + y^2}$ b) $y = \frac{8}{3}x - \frac{1}{3}$
- $y' = \frac{y}{2y - x}$ and $y'' = \frac{-2xy + 2y^2}{(2y - x)^3}$
- The tangent line is horizontal with $x = \pm\sqrt{10}$, $x = \pm\sqrt{2}$, and $x = 0$.
- $x = \frac{-2}{3}$ and $x = -1$
- $g(x)$ has an absolute minimum at $(2, 3)$ and an absolute maximum at $(1, 5)$

14. (a) Both x - and y - intercept at $(0, 0)$.
(b) Vertical asymptote at $x = -\frac{1}{2}$, horizontal asymptote at $y = 0$: $y \rightarrow 0$ as $x \rightarrow \pm\infty$
(c) Increasing on $(-\frac{1}{2}, \frac{1}{2})$ and decreasing on $(-\infty, -\frac{1}{2}) \cup (\frac{1}{2}, \infty)$
(d) Local max at $(\frac{1}{2}, \frac{1}{8})$
(e) Concave down on $(-\infty, -\frac{1}{2}) \cup (-\frac{1}{2}, 1)$ and concave up on $(1, \infty)$
(f) Inflection point at $(1, \frac{1}{9})$.
15. (a) $\overline{C(x)} = \frac{2000}{x} + 12$
(b) $\lim_{x \rightarrow \infty} \overline{C(x)} = 12$, so average cost approaches \$12
(c) Marginal average cost at 300 is $\approx -\$0.02$
16. Maximum area is $464,640m^2$ and is achieved when $x = 88m$ and $y = 528m$.
17. (a) $\eta = \frac{-2p^2}{72081 - p^2}$
(b) Demand will increase by .00675%
(c) Inelastic, so in this case, revenue will increase.