

Question 1: (9 pts) Evaluate the following limits:

$$\text{a) } \lim_{x \rightarrow 0} \frac{e^{2x} + x^2 - 2x - 1}{\cos(3x) - 1} \quad \text{b) } \lim_{x \rightarrow +\infty} (3x - 1)^{\frac{2}{\ln(x)+1}} \quad \text{c) } \lim_{x \rightarrow \pi} \sin(2x) \csc(5x)$$

Question 2: (4 pts) Use Newton's method to find the solution of $\sin(x) = x^2 - 2$ that is between $x = 1$ and $x = 2$. Give an answer that is accurate to 5 decimals. (Show all your work!)

Question 3: (9 pts) Find the derivative of the following functions. **Do not simplify your answer.**

$$\text{a) } y = e^{\arccos(3x)} \quad \text{b) } y = (\arctan(\sqrt{x}))^4 \quad \text{c) } y = \ln(\arcsin(x^3))$$

Question 4: (4 pts) Use the Trapezoidal Rule to approximate $\int_1^2 \sqrt{x^3 - 1} \, dx$, using $n = 5$.

Give an answer that is accurate to 4 decimals.

Question 5: (4 pts) For raising a load, the efficiency of a screw with square threads is given by:

$$E = \frac{100T(1 - fT)}{T + f}$$

where f is the coefficient of friction, and T is the *tangent* of the pitch angle of the screw. If $f = 0.25$, what acute angle will give the maximum efficiency? (Hint: find $\frac{dE}{dT}$)

Question 6: (3 pts) A conveyor belt is dumping gravel at a rate of $40.0 \text{ ft}^3/\text{min}$. The gravel is accumulating in a pile that has a conical shape, and whose height and diameter are always equal.


How fast is the height of the pile increasing when it's 5.00 ft ? (Remember volume of a cone: $V = \frac{\pi}{3}r^2h$)

Question 7: (13 pts)

$$\text{Given } f(x) = \frac{x-1}{x^2-2x}, \quad f'(x) = \frac{-(x^2-2x+2)}{(x^2-2x)^2} \quad \text{and} \quad f''(x) = \frac{2(x-1)(x^2-2x+4)}{(x^2-2x)^3}, \quad \text{find (if any):}$$

- a) The domain of f . b) The x and y intercept(s). c) The vertical and horizontal asymptotes.
d) The intervals on which f is increasing or decreasing. e) The local minima and maxima.
f) The intervals of upward and downward concavity. g) The points of inflection.
h) Sketch the graph of f .

Question 8: (3 pts) Find the area of the region enclosed by the graph of $f(x) = x^3 - x$ and the x -axis:




Question 9: (28 pts) Evaluate the following integrals:

a) $\int \frac{2x^3 - 2x^2 - 6x + 13}{x^2 - x - 2} dx$ b) $\int \tan^3(x) \cos^2(x) dx$ c) $\int \arccos(x) dx$

d) $\int \sqrt{x^2 + 6x + 10} dx$ e) $\int \frac{x^2 + 1}{\sqrt[5]{x^3 + 3x + 7}} dx$ f) $\int \tan^8(3x) \sec^4(3x) dx$

g) $\int (3x^2 + 2) e^{2x} dx$

Question 10: (6 pts) Let R be the region enclosed by $y = 1 + 2x^3$, $y = 0$, $x = 0$ and $x = 1$:



[2] a) Set up, but **do not evaluate**, the integral to find the volume of the solid obtained by revolving R about the line $y = -2$.

[4] b) Find the volume of the solid obtained by revolving R about the y -axis.

Question 11: (3 pts) Determine if $y = x^3 e^{2x}$ is a solution of the differential equation $y'' - 4y' + 6y = 6xe^{2x}$.

Question 12: (4 pts) Solve the following separable differential equation: $2 \sin(x) \frac{dy}{dx} = \frac{\tan(x)}{y}$ with the initial condition $y(0) = -2$.

Question 13: (4 pts) Solve the following first order linear differential equation:

$$x y' = y + \frac{2x^2}{x^2 + 1} \quad \text{with initial condition } y(\pi/4) = \pi$$

Question 14: (6 pts) Given the function $f(x) = \begin{cases} 0 & \text{if } -\pi \leq x < 0 \\ x & \text{if } 0 \leq x < \pi \end{cases}$ find a_0 , a_1 and b_1 of the Fourier series of the function.


ANSWERS:

1.) a) $\frac{-2}{3}$ b) e^2 c) $\frac{-2}{5}$ 2.) 1.72847 3.) a) $e^{\arccos(3x)} \cdot \frac{-1}{\sqrt{1-(3x)^2}} \cdot 3$

b) $4\left(\arctan(\sqrt{x})\right)^3 \cdot \frac{1}{1+(\sqrt{x})^2} \cdot \frac{1}{2\sqrt{x}}$ c) $\frac{1}{\arcsin(x^3)} \cdot \frac{1}{\sqrt{1-(x^3)^2}} \cdot 3x^2$ 4.) 1.4909

5.) 0.6629rad or 37.98° 6.) 2.04 ft/min

- 7.) a) $\mathbb{R} \setminus \{0, 2\}$ b) x-int: (1,0) y-int: none
 c) V.A.: $x = 0$ and $x = 2$ H.A.: $y = 0$
 d) Inc. never, Dec. on $]-\infty, 0[$, $]0, 2[$ and $]2, +\infty[$
 e) None
 f) C.U.: $]0, 1[$ and $]2, +\infty[$ C.D.: $]-\infty, 0[$ and $]1, 2[$ g) (1,0)



8.) $\frac{1}{2}$ 9.) a) $x^2 + 3 \ln|x-2| - 5 \ln|x+1| + C$ b) $\frac{\cos^2(x)}{2} - \ln|\cos(x)| + C$

c) $x \arccos(x) - \sqrt{1-x^2} + C$ d) $\frac{1}{2}(x+3)\sqrt{x^2+6x+10} + \frac{1}{2} \ln|\sqrt{x^2+6x+10} + x+3| + C$

e) $\frac{5}{12}(x^3+3x+7)^{4/5} + C$ f) $\frac{1}{33} \tan^{11}(3x) + \frac{1}{27} \tan^9(3x) + C$

g) $\frac{1}{2}(3x^2+2)e^{2x} - \frac{3}{2}xe^{2x} + \frac{3}{4}e^{2x} + C$ 10.) a) $\int_0^1 \pi[(3+2x^3)^2 - 2^2] dx$ b) $\frac{9\pi}{5}$

11.) It is not a solution 12.) $y = -\sqrt{\ln|\sec(x)\tan(x)| + 4}$ 13.) $y = 2x \arctan(x) + 2x$

14.) $a_0 = \frac{\pi}{4}$ $a_1 = \frac{-2}{\pi}$ $b_1 = 1$